

Impact of prenatal exercise on both prenatal and postnatal anxiety and depressive symptoms: a systematic review and meta-analysis

Margie H Davenport,¹ Ashley P McCurdy,¹ Michelle F Mottola,² Rachel J Skow,¹ Victoria L Meah,³ Veronica J Poitras,⁴ Alejandra Jaramillo Garcia,⁴ Casey E Gray,⁵ Nick Barrowman,⁶ Laurel Riske,¹ Frances Sobierajski,¹ Marina James,¹ Taniya Nagpal,² Andree-Anne Marchand,⁷ Megan Nuspl,⁸ Linda G Slater,⁹ Ruben Barakat,¹⁰ Kristi B Adamo,¹¹ Gregory A Davies,¹² Stephanie-May Ruchat¹³

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/bjsports-2018-099697>).

For numbered affiliations see end of article.

Correspondence to

Dr Margie H Davenport, Program for Pregnancy and Postpartum Health, Faculty of Kinesiology, Sport, and Recreation, University of Alberta, Edmonton T6G 2E1, Canada; mdavenport@ualberta.ca

Accepted 10 August 2018

ABSTRACT

Objective To examine the influence of prenatal exercise on depression and anxiety during pregnancy and the postpartum period.

Design Systematic review with random effects meta-analysis and meta-regression.

Data sources Online databases were searched up to 6 January 2017.

Study eligibility criteria Studies of all designs were included (except case studies) if they were published in English, Spanish or French and contained information on the Population (pregnant women without contraindication to exercise), Intervention (subjective or objective measures of frequency, intensity, duration, volume or type of exercise), Comparator (no exercise or different frequency, intensity, duration, volume and type of exercise) and Outcome (prenatal or postnatal depression or anxiety).

Results A total of 52 studies (n=131 406) were included. 'Moderate' quality evidence from randomised controlled trials (RCTs) revealed that exercise-only interventions, but not exercise+cointerventions, reduced the severity of prenatal depressive symptoms (13 RCTs, n=1076; standardised mean difference: -0.38, 95% CI -0.51 to -0.25, I²=10%) and the odds of prenatal depression by 67% (5 RCTs, n=683; OR: 0.33, 95% CI 0.21 to 0.53, I²=0%) compared with no exercise. Prenatal exercise did not alter the odds of postpartum depression or the severity of depressive symptoms, nor anxiety or anxiety symptoms during or following pregnancy. To achieve at least a moderate effect size in the reduction of the severity of prenatal depressive symptoms, pregnant women needed to accumulate at least 644 MET-min/week of exercise (eg, 150 min of moderate intensity exercise, such as brisk walking, water aerobics, stationary cycling, resistance training).

Summary/Conclusions Prenatal exercise reduced the odds and severity of prenatal depression.

it is generally accepted that both conditions are underdiagnosed and undertreated during the perinatal period.^{5–6} These conditions have immediate and persisting adverse impacts on both mother and child.

For mothers, depression and/or anxiety during pregnancy strongly predicts postpartum depression and anxiety; depression and anxiety are associated with reduced self-care, preterm labour, low birth weight as well as compromised caregiving and bonding with the infant.^{7–9} Maternal depression and anxiety have also been linked with delays in offspring development (cognitive, emotional and social development).^{10–13} Standard treatments for depression and anxiety include pharmacological and psychological intervention^{14–16} but the long-term impact of some medications on the fetus has not been established^{7–16} and psychotherapy can be costly and difficult to access.^{17–18} The consequences of leaving these disorders untreated can be severe.

In non-pregnant women, exercise may be as effective as antidepressants and psychotherapy for treating mild-to-moderate depression (ie, a moderate effect size).¹⁹ A recent review demonstrated that light to moderate intensity aerobic exercise initiated in the postpartum period improves mild to moderate depressive symptoms and increases the likelihood of resolving mild to moderate depression in the postpartum period compared with standard care.²⁰ It has not been established if prenatal exercise reduces the prevalence and/or severity of depression and anxiety during pregnancy and the postpartum period.

Current national and international guidelines recommend that women without contraindications to exercise (eg, preeclampsia, incompetent cervix, premature labour) be physically active throughout pregnancy.^{21–23} This systematic review and meta-analysis forms part of a series of reviews, which will form the evidence base for the development of the 2019 *Canadian guideline for physical activity throughout pregnancy* (herein referred to as *Guideline*).²⁴ The purpose of this paper is to present the results of a systematic review and meta-analysis of the relationships between prenatal exercise, and (1) depression and anxiety during pregnancy as well as (2) any persisting effect into the postpartum period.

INTRODUCTION

In 2017, the WHO identified that depression is the leading cause of disability worldwide.¹ Pregnant and postpartum women are particularly vulnerable to declining mental health as a result of elevated stress and lack of social support.² Depression affects approximately 13% and anxiety affects up to 39% of pregnant and postpartum women;^{3–4} however,



© Author(s) (or their employer(s)) 2018. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Davenport MH, McCurdy AP, Mottola MF, et al. *Br J Sports Med* 2018;**52**:1376–1385.

METHODS

In October 2015, the Guidelines Consensus Panel was formed to identify outcomes for the *Guideline* update. The panel included researchers, methodological experts, a fitness professional, a public health representative (Middlesex-London Health Unit) and representatives from the Society for Obstetricians and Gynecologists of Canada, Canadian Society for Exercise Physiology (CSEP), The College of Family Physicians of Canada, Canadian Association of Midwives and Canadian Academy of Sport and Exercise Medicine, Exercise is Medicine Canada. During this meeting, 20 ‘critical’ and 17 ‘important’ outcomes related to prenatal exercise and maternal/fetal health were selected. Maternal mental health was rated as ‘critical’ outcome. This systematic review and meta-analysis was conducted in accordance with the PRISMA guidelines and the checklist was completed.²⁵

Protocol and registration

Two systematic reviews examining the impact of prenatal exercise on fetal and maternal health outcomes were registered with PROSPERO, the international prospective register of systematic reviews (*fetal health*: Registration no. CRD42016029869; Available from: https://www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42016029869; *maternal health*: Registration no. CRD42016032376; Available from: https://www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42016032376). Because the relationships between prenatal exercise and maternal/fetal health outcomes are examined in studies related to both fetal and maternal health, records retrieved from both of these reviews were considered for inclusion in the present systematic review.

Eligibility criteria

This study was guided by the participants, interventions, comparisons, outcomes and study design (PICOS) framework.²⁵

POPULATION

The population of interest included in this review was pregnant women without contraindication to exercise (according to the CSEP and American College of Obstetricians and Gynecologists (ACOG) guidelines).^{21 26} Absolute contraindications to exercise were defined as: ruptured membranes, premature labour, persistent second or third trimester bleeding, placenta praevia, preeclampsia, gestational hypertension, incompetent cervix, intrauterine growth restriction, high order pregnancy, uncontrolled type 1 diabetes, hypertension or thyroid disease or other serious cardiovascular, respiratory or systemic disorders. Relative contraindications to exercise were defined as: a history of spontaneous abortion, premature labour mild/moderate cardiovascular or respiratory disease, anaemia or iron deficiency, malnutrition or eating disorder, twin pregnancy after 28 weeks or other significant medical conditions.^{21 22 26}

INTERVENTION (EXPOSURE)

The intervention/exposure was subjective or objective measures of frequency, intensity, duration, volume or type of exercise. Although exercise is a subtype of physical activity, for the purpose of this review, we used the terms interchangeably. Exercise and physical activity were defined as any bodily movement generated by skeletal muscles that resulted in energy expenditure above resting levels.²⁷ Acute (ie, a single exercise session) or habitual (ie, usual activity) prenatal exercise and interventions including exercise alone (termed ‘exercise-only’ interventions)

or in combination with other interventions (such as diet; termed ‘exercise+cointerventions’) were considered. Studies were excluded if exercise was performed after the beginning of labour.

COMPARISON

Eligible comparators were: various frequencies, intensities, durations, volumes and types of exercise; various durations of intervention or exercise or exercise in different trimesters.

OUTCOME

Eligible outcomes were diagnosis of depression or anxiety (either by exceeding a specific threshold via questionnaire or diagnosis by a qualified professional) and severity of symptoms of depression or anxiety.

STUDY DESIGN

Primary studies of any design were eligible, except case studies (n=1). Narrative or systematic reviews and meta-analyses were excluded.

Information sources

A comprehensive search was created and run by a research librarian (LGS) in the following databases: MEDLINE, EMBASE, PsycINFO, Cochrane Database of Systematic Reviews, Cochrane Central Register of Controlled Trials, Scopus and Web of Science Core Collection, CINAHL Plus with Full-text, Child Development & Adolescent Studies, ERIC, Sport Discus, ClinicalTrials.gov and the Trip Database up to 6 January 2017 (see online supplementary file for complete search strategies).

Study selection and data extraction

Titles and abstracts of all retrieved articles were independently screened by two reviewers. Abstracts that were judged to have met the initial screening criteria by at least one reviewer were retrieved as full-text articles. Articles were reviewed for relevant PICOS information by at least one person. If it was deemed that articles did not meet the inclusion criteria, they were reviewed by MHD and/or SMR prior to exclusion. If agreement could not be reached by discussion, study characteristics related to eligibility were presented to the Guidelines Steering Committee (MHD, MFM, SMR, CG, VP, AJG and NB) and final decisions were made by consensus. Studies that were selected were imported into DistillerSR (Evidence Partners, Ottawa, Ontario, Canada) for data extraction. At this point, studies from the maternal and fetal reviews that were included were de-duplicated against one another in DistillerSR and were considered as one review from this point forward.

Data extraction tables were created in DistillerSR in consultation with methodological experts and the Guidelines Steering Committee. Data were extracted by one person; a content expert (MHD, MFM or SMR) independently verified the extracted data. Reviewers were not blinded to study authors. For each single study, the most recent or complete version of the publication was selected as the ‘parent’ paper; however, relevant data from all publications related to each unique study were extracted. Study characteristics (ie, year, study design, country) and population characteristics (eg, number of participants, age, pre-pregnancy body mass index (BMI), previous physical activity levels, parity and pregnancy complications including preeclampsia, gestational hypertension and gestational diabetes), intervention/exposure (actual and/or prescribed exercise frequency, intensity, duration and type, duration of the intervention, measure of physical activity) and outcomes (depression or anxiety diagnosis and/or

symptom severity) were extracted (see online supplementary table 1). If data were not available for extraction, the authors were contacted for additional information.

Quality of evidence assessment

The Grading of Recommendations Assessment, Development and Evaluation (GRADE) framework was used to assess the quality of evidence across studies for each study design and health outcome. The risk of bias for all included studies was independently assessed by two reviewers. The risk of bias in randomised controlled trials (RCTs) and intervention studies was evaluated following the Cochrane Handbook²⁸; risk of bias in observational studies was assessed using the characteristics recommended by Guyatt *et al.*²⁹ All studies were examined for potential sources of bias, ie, selection bias, reporting bias, performance bias, detection bias, attrition bias and 'other' sources of bias.

Risk of bias across studies was rated as 'serious' when studies having the greatest influence on the pooled result (assessed using weight (%) given in forest plots or sample size in studies that were narratively synthesised) presented 'high' risk of bias. The greatest influence on the pooled result was determined as follows: the studies that had the greatest individual per cent contribution in the meta-analyses, when taken together, contributed to >50% of the weight of the pooled estimate. Serious risk of bias was considered when sample size of studies that were narratively synthesised was similar to the total sample size of studies contributing to >50% of the weight of the pooled estimate in the meta-analyses. Performance bias was rated as 'high' when <60% of participants performed 100% of prescribed exercise sessions or attended 100% of counselling sessions (defined as low compliance) or when compliance to the intervention was not reported. Attrition bias was rated as 'high' when >10% of data were missing at the end of the study and intention-to-treat analysis was not used. Given the nature of exercise interventions, it is not possible to blind participants to group allocation. Therefore, if the only source of bias was related to the blinding of allocation, the risk of bias was rated as 'low'.

The risk of bias, indirectness, inconsistency, imprecision or risk of publication bias was assessed across studies for each study design and health outcome. Evidence from RCTs began with a 'high' quality of evidence rating and was graded down if any quality violations occurred. Evidence from all non-randomised intervention and observational studies began with a 'low' quality rating and, if there was no cause to downgrade, was upgraded if applicable according to the GRADE criteria (eg, large magnitude of effect, evidence of dose-response).²⁸

Indirectness was considered serious when exercise-only interventions and exercise+cointerventions were combined for analysis or when the effect of exercise+cointervention on an outcome was examined. Exercise-only interventions could include standard care. Inconsistency was considered serious when heterogeneity was high ($I^2 \geq 50\%$) or when only one study was assessed (I^2 unavailable). Imprecision was considered serious when the 95%CI crossed the line of no effect and was wide, such that interpretation of the data would be different if the true effect were at one end of the CI or the other. When only one study was included, imprecision was not considered serious because inconsistency was already considered serious for this reason. Finally, in order to assess publication bias, funnel plots were created if at least 10 studies were included in the forest plot (see online supplementary figure 2). If there were fewer than 10 studies, publication bias was deemed non-estimable and not rated down. Due to time constraints and feasibility, one reviewer evaluated the quality of the evidence across each health

outcome using the protocol and a second person reviewed the GRADE tables as a quality control measure. Quality of evidence assessment is presented in online supplementary tables 2–5.

Statistical analysis

Statistical analyses were conducted using Review Manager V5.3. (Cochrane Collaboration, Copenhagen, Denmark). ORs were calculated for all dichotomous outcomes. Inverse-variance weighting was applied to obtain OR using a random effects model. Standardised mean differences (SMD) were calculated when different measurement tools or scales were used for a single outcome. SMD effect sizes were calculated using Hedges' g. An effect size of 0.2, 0.4 and 0.8 was considered small, moderate and large, respectively. Significance was defined as $p < 0.05$. Meta-analyses were performed separately by study design. A staged approach was used to determine if there was sufficient evidence from RCTs for each outcome to inform the *Guideline* or if it was necessary to consider other study designs. If fewer than 2000 women were included in the exercise-only RCT meta-analysis, the impact of prenatal exercise on the specific outcome was examined further using observational evidence (non-randomised intervention, cohort, cross-sectional and case-control studies). For RCTs and non-randomised interventions, sensitivity analyses were performed to evaluate whether the effects were different when examining relationships between exercise-only interventions versus exercise+cointerventions and depression and anxiety. When possible, the following a priori subgroup analyses were conducted for exercise-only interventions: (1) women diagnosed with diabetes (gestational, type 1 or type 2) compared with women without diabetes (named 'general population'); (2) samples of women with overweight or obesity (mean BMI > 25.0 kg/m²) prior to pregnancy compared with samples of women who were of various BMI (mean BMI < 25 kg/m² but possibly with some individuals with BMI > 25.0 kg/m²; named 'general population'); (3) women > 35 years of age compared with women < 35 years of age; (4) women who were previously inactive compared with those who were previously active (as defined by individual study authors). If a study did not provide sufficient detail to allow it to be grouped into the a priori subgroups, then a third group called 'unspecified' was created. Tests for subgroup differences were conducted, with statistical significance set at $p < 0.05$. If statistically significant differences were found subgroup differences were interpreted. Finally, a priori subgroup analyses were also conducted for exercise-only RCTs with critical outcomes to identify whether a specific type of exercise was associated with greater benefit compared with aerobic or resistance training alone. *Posthoc* subgroup analyses were conducted comparing women with and without depression at baseline, women who exercised in supervised versus unsupervised settings and comparing effects in which the Edinburgh Postnatal Depression Scale (EPDS) versus Center for Epidemiologic Studies Depression Scale (CES-D) tools were used to assess possible diagnosis of depression and symptom severity. The I^2 was calculated to indicate the per cent of total variability that was attributable to between-study heterogeneity. In studies where there were no observed events in the intervention or control groups, data were entered into forest plots, but were considered 'not estimable' and excluded from the pooled analysis as per the recommendation in the Cochrane Handbook.³⁰ Dose-response meta-regression^{31 32} was carried out by weighted no-intercept regression of log OR with a random effects for study, using the metafor³³ package in R³⁴ V3.4.1. Models did not include an intercept term since the log OR is assumed to be zero when the exercise dose is zero. Restricted cubic splines with knots at the 10th, 50th and 90th percentiles of the explanatory variable³⁵ were

used to investigate whether there was evidence for a nonlinear relationship. Fitting was performed by maximum likelihood, and nonlinearity was assessed using a likelihood ratio test. When the model was statistically significant at $p < 0.05$, the minimum exercise dose to obtain a clinically meaningful benefit was estimated by the minimum value of the explanatory variable at which the estimated OR was less than 0.75. Linear models were presented unless the fit of the spline was significantly better. For outcomes or for subsets of studies where a meta-analysis was not possible, a narrative synthesis of the results was conducted, organised around each outcome. Unless otherwise specified, studies were not included in meta-analyses if data were reported incompletely (eg, SD, SE or number of cases/controls not provided), if data were adjusted for confounding factors or if the study did not include a non-exercising control group. In studies where data were included in the meta-analysis but additional information was available that could not be meta-analysed, the studies were incorporated in both the

meta-analysis and narrative synthesis. Within each outcome, results were presented by study design.

RESULTS

Study selection

The initial search was not limited by language. However, the Guidelines Steering Committee decided to exclude studies published in languages other than English, Spanish or French for feasibility reasons. A PRISMA diagram of the search results, including reasons for exclusion, is shown in figure 1. A comprehensive list of excluded studies is presented in the online supplementary file.

Study characteristics

Overall, 52 unique studies ($n = 131\,406$ women) from 19 countries were included. There were 26 RCTs, 7 non-randomised intervention, 10 cohort, 6 cross-sectional and 3 case control




Figure 1 Flow diagram of studies selected for the present study.

studies. Among the included exercise interventions, the frequency of exercise ranged from 1 to 7 days per week, the duration of exercise ranged from 20 to 75 min per session and the types of exercise included aerobic exercise, yoga, resistance training and pelvic floor muscle training. Additional details about the studies can be found in the online supplementary file. Possible diagnosis of depression or being high in anxiety was based on a cut-off score on a clinical scale. Severity of symptoms was the overall score on a clinical scale. State anxiety is an acute response to a perceived or real threat, while trait anxiety describes the inter-individual tendency to have high state anxiety in response to a threat.

Quality of evidence

Overall, the quality of evidence ranged from 'very low' to 'high' (see online supplementary tables 2 and 3). The most common reasons for downgrading the quality of evidence were (1) serious risk of bias that reduced the level of confidence in the observed effects and (2) indirectness of the interventions being assessed. Common sources of bias included poor or unreported compliance and inappropriate treatment of missing data when attrition rate was high. Publication bias was not observed among the analyses where it was possible to systematically assess publication bias using funnel plots.

Synthesis of data

The results of the meta-regression analysis are presented in the online supplementary file (*Meta-regressions*, online supplementary figures 31–34). Meta-regression analysis using linear and spline regression was conducted for each outcome. In each case, the spline model did not provide a significantly better fit

than the linear model. Minimum exercise thresholds required to achieve a moderate effect size ($SMD > 0.4$) were estimated. In order to achieve a moderate effect of exercise in reducing prenatal depressive symptoms, the following thresholds were identified: an exercise intensity of 4.0 METs/min (eg, light walking; online supplementary figure 31), exercise duration > 49.9 min per session (online supplementary figure 32); exercise frequency > 2.6 times per week (online supplementary figure 33) and volume of exercise per week > 644 MET-min/week (eg, 150 min of moderate intensity exercise, such as brisk walking, water aerobics, stationary cycling, resistance training; online supplementary figure 34).

Prenatal depressive symptoms

Overall, there was 'very low' quality evidence from 21 RCTs ($n=4267$) regarding the association between prenatal exercise and depressive symptoms. The quality of evidence was downgraded from 'high' to 'very low' because of serious risk of bias, serious inconsistency and serious indirectness of the intervention. Overall, prenatal exercise was associated with a small reduction in severity of depressive symptoms compared with no exercise (pooled estimate based on 19 RCTs, $n=3316$; $SMD: -0.23$, 95% CI -0.36 to -0.09 , $I^2=63\%$; figure 2). Two exercise-only interventions could not be included in the meta-analysis^{36 37} (see online supplementary file). In the first RCT, there was no influence of prenatal exercise on the severity of depressive symptoms during pregnancy.³⁷ In contrast, in the second RCT, yoga ($n=51$) but not non-yoga ($n=45$) antenatal exercises was associated with an improvement in depressive symptoms.³⁶




Figure 2 Effects of prenatal exercise-only versus exercise plus co-interventions (randomised controlled trials) compared with control on prenatal depressive symptoms. Analyses conducted with a random effects model. IV, inverse variance.

Sensitivity analysis

The pooled estimate for the exercise-only interventions was significantly different than the exercise+cointervention subgroups ($p < 0.0001$). Specifically, exercise-only interventions had a small effect on reducing the severity of depressive symptoms compared with no exercise (13 RCTs, $n = 1076$; SMD: -0.38 , 95% CI -0.51 to -0.25 , $I^2 = 10\%$; 'moderate' quality evidence, downgraded due to serious risk of bias; online supplementary figure 1). There was no statistically significant difference for those participating in exercise+cointerventions (online supplementary figure 1).

Subgroup analyses

There were no statistically significant subgroup differences identified (see figure 2, online supplementary figure 3–7).

Other study designs

Findings from four non-randomised interventions,^{38–41} one cohort⁴² and three cross-sectional studies^{43–45} were in agreement with the findings from RCTs. In contrast, findings from three cohort studies^{46–48} and one case-control study⁴⁹ were not significant (see online supplementary file for more details).

Prenatal depression

Overall, there was 'very low' quality evidence from eight RCTs ($n = 2481$) regarding the association between prenatal exercise and prenatal depression. The quality of evidence was downgraded from 'high' to 'very low' because of serious risk of bias, serious inconsistency and serious indirectness of the intervention. Overall, prenatal exercise was associated with 45% lower odds of developing prenatal depression compared with no exercise (OR 0.55, 95% CI 0.34 to 0.90, $I^2 = 59\%$; figure 3).

Sensitivity analysis

The pooled estimate for the exercise-only interventions was significantly different than the exercise+cointervention subgroups ($p < 0.0001$). Specifically, exercise-only interventions (all interventions included supervised exercise) reduced the odds of having depression by 67% (5 RCTs, $n = 683$; OR: 0.33, 95% CI 0.21 to 0.53, $I^2 = 0\%$; 'moderate' quality evidence, downgraded due to serious risk of bias; figure 3). There was no statistically significant difference in odds of developing depression for those participating in exercise+cointerventions compared with a no-exercise control (figure 3).

Subgroup analyses

Stratification by exercise type showed no statistically significant difference in odds of developing prenatal depression (see online supplementary figure 13). The remaining a priori subgroup analyses could not be conducted as studies with the relevant subgroups did not exist.

Other study designs

Findings from one non-randomised intervention,³⁹ one cohort study⁴⁷ and one cross-sectional study⁵⁰ were in agreement with the findings from RCTs (see online supplementary file for more details).

Postnatal depressive symptoms

Overall, there was 'low' quality evidence from seven RCTs ($n = 2795$) regarding the association between prenatal exercise and severity of postnatal depressive symptoms. The quality of evidence was downgraded from 'high' to 'very low' because of serious risk of bias and serious indirectness of the intervention. Overall, prenatal exercise was not associated with reduction in postnatal depressive symptoms compared with no exercise (SMD: 0.05, 95% CI -0.02 to 0.12 , $I^2 = 0\%$; figure 4).




Figure 3 Effects of prenatal exercise-only versus exercise+cointervention compared with control on prenatal depression (randomised controlled trials). Analyses conducted with a random effects model. M-H, Mantel-Haenszel method.




Figure 4 Effects of prenatal exercise-only versus exercise+cointervention compared with control on postnatal depressive symptoms (randomised controlled trials). Analyses conducted with a random effects model. IV, inverse variance.

Sensitivity analysis

The pooled estimate for the exercise-only interventions was not significantly different than the exercise+cointervention subgroups ($p=0.26$). Prenatal exercise-only interventions did not affect the severity of depressive symptoms in the postpartum period (figure 4).

Subgroup analyses

Stratification by exercise type showed no statistically significant difference in odds of developing postnatal depressive symptoms. The remaining subgroup analyses on exercise-only intervention could not be conducted due to a lack of studies with relevant subgroups.

Other study designs

The findings from three non-randomised interventions,^{38 40 51} one cohort study⁴⁶ and one cross-sectional study⁵² are presented in the online supplementary file.

Postnatal depression

Overall, there was 'very low' quality evidence from four RCTs ($n=2536$) regarding the association between prenatal exercise and postnatal depression. The quality of evidence was downgraded from 'high' to 'very low' because of serious risk of bias, serious imprecision and serious indirectness of the intervention. Overall, prenatal exercise did not decrease the odds of having depression in the postpartum period compared with no exercise (OR 0.91, 95% CI 0.61 to 1.36, $I^2=8\%$; online supplementary figure 18).

Sensitivity analysis

The pooled estimate for the exercise-only interventions was not significantly different than the pooled estimate for the exercise+cointervention subgroups ($p=0.12$). Prenatal exercise-only interventions did not affect the odds of depression in the postpartum period (online supplementary figure 18).

Subgroup analyses

Stratification by exercise type indicated no statistically significant difference in odds of developing postnatal depression (see online supplementary figure 19). The remaining subgroup analyses on exercise-only intervention could not be conducted due to a lack of studies with relevant subgroups.

Other study designs

There was 'very low' quality evidence (downgraded due to serious risk of bias and serious inconsistency) from one cohort study ($n=1305$)⁵³ showing a reduced odds of postnatal with prenatal exercise (OR 0.79, 95% CI 0.70 to 0.89; online supplementary figure 20).

Prenatal state anxiety symptoms

Overall, there was 'very low' quality evidence from eight RCTs ($n=1785$) regarding the association between prenatal exercise and prenatal state anxiety symptoms.³⁶ The quality of evidence was downgraded from 'high' to 'very low' because of serious risk of bias, serious imprecision and serious indirectness of the intervention. Overall, prenatal exercise was not associated with a reduction in prenatal state anxiety symptoms compared with no exercise (pooled estimate based on seven RCTs, $n=1689$; SMD: 0.06, 95% CI -0.04 to 0.15, $I^2=0\%$; online supplementary figure 21). The one superiority trial that could not be included in the meta-analysis showed an improvement in state anxiety symptoms with yoga ($n=51$), but not with other forms of antenatal exercise ($n=45$).³⁶

Sensitivity analyses

The pooled estimate for the exercise-only interventions was not significantly different than the pooled estimate for exercise+cointervention subgroups ($p=0.79$). Prenatal exercise-only interventions did not affect the severity of prenatal state anxiety symptoms during pregnancy in either group (online supplementary figure 21).




Figure 5 Effects of prenatal exercise compared with control on prenatal state anxiety symptoms (randomised controlled trials). Subgroup analyses were conducted with studies including women with depression prior to the intervention with women who without depression (defined as 'general population'). Analyses conducted with a random effects model. IV, inverse variance.

Subgroup analyses

The tests for subgroup differences performed for exercise-only interventions were not statistically significant (figure 5).

Other study designs

The findings from one non-randomised intervention⁵⁴ and one cohort study⁴⁷ were in agreement with the findings from RCT. In contrast, one cross-sectional study³⁸ did not report a reduction in prenatal state anxiety symptoms with prenatal exercise. See online supplementary file for more details.

Prenatal state anxiety

Overall, there was 'very low' quality evidence from two RCTs of exercise+cointervention ($n=1534$)^{55,56} regarding the association between prenatal exercise and odds of prenatal state anxiety. The quality of evidence was downgraded from 'high' to 'very low' because of serious risk of bias, serious indirectness of the intervention and serious imprecision. Overall, prenatal exercise was not associated with lower odds of prenatal state anxiety compared with no exercise (OR: 1.12, 95% CI 0.85 to 1.48, $I^2=0\%$; online supplementary figure 22).

Sensitivity analysis

Sensitivity analysis could not be conducted as there were no exercise-only interventions.

Prenatal trait anxiety symptoms

Overall, there was 'low' quality evidence from two RCTs of exercise-only interventions ($n=90$)⁵⁷ regarding the association between prenatal exercise and severity of prenatal trait anxiety symptoms. The quality of evidence was downgraded from 'high' to 'low' because of serious risk of bias and serious imprecision. Overall, prenatal exercise was not associated with a decrease in prenatal trait anxiety symptoms compared with no exercise (SMD: -0.21, 95% CI -0.63 to 0.20, $I^2=0\%$; online supplementary figure 25). The one superiority trial that could not be included in the meta-analysis showed an improvement in trait anxiety symptoms with yoga ($n=51$), but not with other antenatal exercise ($n=45$).³⁶

Sensitivity analysis

Sensitivity analysis could not be conducted as there were no exercise+cointervention.

Other study designs

Findings from one non-randomised intervention⁵⁸ and one case control study⁴⁹ were in agreement with findings from RCTs (online supplementary figures 26 and 27), while a cross-sectional⁵⁹ study demonstrated a reduction in the severity of prenatal trait anxiety symptoms in women who exercised during pregnancy compared with those who were inactive ($n=203$; SMD: -0.73, 95% CI -1.02 to -0.45; online supplementary figure 28).

Prenatal trait anxiety

Overall, there was 'very low' quality evidence from one RCT of exercise+cointerventions ($n=161$) regarding the association between prenatal exercise and prenatal trait anxiety.⁵⁵ The quality of evidence was downgraded from 'high' to 'very low' because of serious risk of bias, serious indirectness and serious inconsistency. Overall, prenatal exercise was not associated with lower odds of prenatal trait anxiety compared with no exercise (OR: 0.75, 95% CI 0.36 to 1.56; online supplementary figure 29).

Postnatal state anxiety symptoms

Overall, there was 'very low' quality evidence from two RCTs ($n=1299$)^{56,60} regarding the association between prenatal exercise and postpartum state anxiety symptoms. The quality of evidence was downgraded from 'high' to 'very low' because of serious risk of bias, serious indirectness of the intervention and serious imprecision. Overall, prenatal exercise was not associated with reduction in postpartum state anxiety symptoms compared with no exercise (SMD: 0.01, 95% CI -0.10 to 0.12, $I^2=0\%$; online supplementary figure 30).

Sensitivity analysis

The pooled estimate for the exercise-only interventions was not significantly different than the pooled estimate for the exercise+cointervention subgroups ($p=0.99$). Prenatal

exercise-only interventions did not affect the severity of postnatal state anxiety symptoms (online supplementary figure 30).

Postnatal state anxiety

Overall, there was 'very low' quality evidence from one exercise+cointervention RCT (n=1220)⁵⁶ regarding the association between prenatal exercise and postnatal state anxiety. The quality of evidence was downgraded from 'high' to 'very low' because of serious risk of bias, serious indirectness and serious inconsistency. Overall, prenatal exercise was not associated with lower odds of postnatal state anxiety compared with no exercise (OR: 1.28, 95% CI 0.91 to 1.80; online supplementary figure 31).

DISCUSSION

The main finding of our systematic review was that exercise-only interventions were associated with a reduction in the severity of prenatal depressive symptoms (SMD -0.39), and this reduction was greatest in women who were supervised during exercise. Pregnant women need to accumulate at least 644 MET-min/week of moderate-to-vigorous intensity exercise (eg, 150 min of moderate intensity exercise, such as brisk walking, water aerobics, stationary cycling, resistance training) in order to achieve a moderate effect size in the reduction in the severity of prenatal depressive symptoms. There appears to be a dose-dependent association of this benefit. A greater volume of exercise was associated with a greater reduction in the severity of prenatal depressive symptoms. Further, exercise-only interventions were associated with a 67% reduction in the odds of developing prenatal depression.

Postnatal depression—symptoms and diagnosis—was not reduced by prenatal exercise. There was no association between prenatal exercise and anxiety or the severity of anxiety symptoms in the prenatal or postnatal period.

SENSITIVITY ANALYSIS

Sensitivity analyses comparing trials with cointerventions (diet+exercise; education classes+exercise; smoking cessation programme+exercise) against trials without cointerventions lowered heterogeneity within each group (11% and 0%, respectively). There was a significant difference between the two types of interventions; severity of depressive symptoms was lower postintervention in standalone exercise trials than for trials including cointerventions. The lack of improvement in depressive symptoms in response to a co-intervention may have been due to the fact that women were required to change multiple behaviours at once, in addition to responding to the demands of pregnancy.⁶¹ Thus, setting realistic and incremental goals may be important considerations when prescribing exercise in the treatment of depression.⁶²

Supervised exercise was associated with a moderate reduction in the severity of depressive symptoms. This may be partially explained by the social support women in supervised interventions received; low social support is a known correlate of depression.¹⁶ It has been suggested that supervision enhances compliance and effectiveness of interventions.⁶³ However, because adherence was either not reported by the authors in a standard manner or not reported at all, we were not able to analyse the effects of the exercise-only interventions in compliant versus non-compliant women to focus on the issue of compliance.

What mechanisms may underpin these epidemiological findings?

Rauff and Downs suggested that improvement in body image may be a factor mediating the positive relationship between exercise and reduced depression or depressive symptoms during pregnancy.⁶⁴ Physiological mechanisms have also been proposed. In non-pregnant populations, aerobic exercise may normalise dopamine-serotonin levels and reduce stress, resulting in a subsequent decrease in anxiety and depression.^{65 66} Depression has been suggested to result, at least in part, from dysregulation of biochemical and neurophysiological function including attenuated release of norepinephrine, serotonin and dopamine.⁶⁷ Exercise may counteract these effects by increasing the release of dopamine and serotonin as well as blunting the release of cortisol in response to stress.⁶⁸ In our systematic review of exercise-only interventions, both yoga and aerobic exercise had a small effect size in reducing the severity of depressive symptoms. However, these data were from a small number of trials and women; thus, additional studies examining the potential impact of different types of exercise are warranted.

Is there a carryover benefit into the postpartum period?

Prenatal exercise-only interventions had no impact on depressive symptoms or the incidence of depression during the postpartum period. However, a recent meta-analysis suggested that exercise interventions initiated during the postpartum period (at least 4 weeks after delivery) reduced the severity of depressive symptoms.²⁰

We highlight that no trials examined populations with diagnosed clinical anxiety disorders. Our meta-analysis does not extend to the efficacy of exercise in the treatment of anxiety disorders (generalised anxiety disorder, panic disorder, social anxiety disorder, among others).

Strengths of this review include our broad inclusion criteria covering all studies types (except case studies), grey literature and three languages (English, French and Spanish from 19 countries). We applied rigorous methodology (GRADE, PRISMA) to the systematic review process and evaluation of the quality of evidence. Subgroup analyses allowed identification of possible sources of heterogeneity; however, statistical heterogeneity was still moderate within subgroups. There was also considerable variability regarding the tools used to measure depression/depressive symptoms, with the majority of trials using either the CES-D or EPDS. As noted by Gong (2015), the CES-D, while a validated measure of perinatal depression, may misattribute some of the somatic symptoms associated with pregnancy (lack of energy, tiredness) to depression, which may have underestimated the effectiveness of exercise interventions. Five trials examined prenatal exercise and depression or anxiety and showed considerable baseline differences in depression and anxiety. However, an exploratory sensitivity analysis performed without these trials had no significant impact on findings. Finally, some of the trials included in our review included women who were using antidepressants, an important confounding variable, that may have influenced the findings.

CONCLUSION

Overall, prenatal exercise reduced the odds of prenatal depression and the severity of depressive symptoms. This positive effect of prenatal exercise did not extend to the postpartum period or prenatal or postnatal anxiety.

What are the new findings?

- ▶ Exercise-only interventions reduced the severity of prenatal depressive symptoms with an effect size similar to that found for psychological treatments in depressed prenatal women.
- ▶ Odds of prenatal depression were reduced by 67% in women who engaged in prenatal exercise.
- ▶ The benefits of prenatal exercise on depression or depressive symptoms did not extend into the postpartum period.
- ▶ To achieve at least a moderate reduction in the severity of prenatal depressive symptoms, pregnant women needed to accumulate at least 644 MET-min/week of exercise (eg, 150 min of moderate intensity exercise, such as brisk walking, water aerobics, stationary cycling, resistance training).

What is already known?

- ▶ Depression and anxiety during and following pregnancy impact the short-term and long-term health issues for mother and child.
- ▶ Exercise prevents and treat anxiety and depression in non-pregnant populations. Whether exercise prevents and treats mental health issues during pregnancy is poorly understood.

Author affiliations

¹Program for Pregnancy and Postpartum Health, Physical Activity and Diabetes Laboratory, Faculty of Kinesiology, Sport, and Recreation, Women and Children's Health Research Institute, Alberta Diabetes Institute, University of Alberta, Edmonton, Alberta, Canada

²R. Samuel McLaughlin Foundation-Exercise and Pregnancy Laboratory, School of Kinesiology, Faculty of Health Sciences, Department of Anatomy and Cell Biology, Schulich School of Medicine and Dentistry, Children's Health Research Institute, The University of Western Ontario, London, Ontario, Canada

³Cardiff School of Sport and Health Sciences, Cardiff Metropolitan University, Cardiff, UK

⁴Independent Researcher, Ottawa, Ontario, Canada

⁵Healthy Active Living and Obesity Research Group, Children's Hospital of Eastern Ontario Research Institute, Ottawa, Ontario, Canada

⁶Clinical Research Unit, Children's Hospital of Eastern Ontario Research Institute, Ottawa, Ontario, Canada

⁷Department of Anatomy, Université du Québec à Trois-Rivières, Trois-Rivières, Québec, Canada

⁸Alberta Research Centre for Health Evidence, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada

⁹John W. Scott Health Sciences Library, University of Alberta, Edmonton, Alberta, Canada

¹⁰Facultad de Ciencias de la Actividad Física y del Deporte-INEF, Universidad Politécnica de Madrid, Madrid, Spain

¹¹School of Human Kinetics, Faculty of Health Sciences, University of Ottawa, Ottawa, Ontario, Canada

¹²Department of Obstetrics and Gynecology, Queen's University, Kingston, Ontario, Canada

¹³Department of Human Kinetics, Université du Québec à Trois-Rivières, Trois-Rivières, Québec, Canada

Acknowledgements We would like to thank Bailey Shandro (UAlberta), Anne Courbalay (UQTR) and Meghan Sebastianski (Alberta SPOR SUPPORT Unit Knowledge Translation Platform), University of Alberta for their assistance with the meta-analysis. We also thank Dr Linda May for her critical review of the manuscript. The authors wish to acknowledge Mary Duggan from the Canadian Society for Exercise Physiology which is one of the primary knowledge users.

Contributors MHD, S-MR, MFM, GAD and KBA contributed to the conception of the study. MHD, S-MR, MFM, GAD, KBA, AJG, NB, VJP, CEG, LGS and RB contributed to the design of the study and development of the search strategy. LGS conducted the systematic search. APM, RS, VLM, LR, FS, MJ, TN, A-AM and MN completed the acquisition of data. MHD and NB performed the data analysis. All authors assisted with the interpretation. MHD and APM were the principal writers of the manuscript. All authors contributed to the drafting and revision of the final article. All authors approved the final submitted version of the manuscript.

Funding This project was funded by a Canadian Institute of Health Research Knowledge Synthesis Grant (140995). MHD is funded by an Advancing Women's Heart Health Initiative New Investigator Award supported by Health Canada and the Heart and Stroke Foundation of Canada (RES0033140). RS is funded by a Canadian Institutes for Health Research Doctoral Research Award (146252). A-AM is funded by a Fonds de Recherche du Québec-Santé Doctoral Research Award (34399).

Competing interests None declared.

Patient consent Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Friedrich MJ. Depression is the leading cause of disability around the world. *JAMA* 2017;317:1517.
- Biaggi A, Conroy S, Pawlby S, *et al.* Identifying the women at risk of antenatal anxiety and depression: A systematic review. *J Affect Disord* 2016;191:62–77.
- Gaynes BN, Gavin N, Meltzer-Brody S, *et al.* Perinatal depression: prevalence, screening accuracy, and screening outcomes. *Evid Rep Technol Assess* 2005;119:1–8.
- Leach LS, Poyser C, Fairweather-Schmidt K. Maternal perinatal anxiety: A review of prevalence and correlates. *Clin Psychol* 2017;21:4–19.
- Goodman JH, Chenauskyl KL, Freeman MP. Anxiety disorders during pregnancy: a systematic review. *J Clin Psychiatry* 2014;75:e1153–e1184.
- Grigoriadis S, VonderPorten EH, Mamisashvili L, *et al.* The impact of maternal depression during pregnancy on perinatal outcomes: a systematic review and meta-analysis. *J Clin Psychiatry* 2013;74:e321–e341.
- Field T. Postpartum depression effects on early interactions, parenting, and safety practices: a review. *Infant Behav Dev* 2010;33:1–6.
- Arteche A, Joormann J, Harvey A, *et al.* The effects of postnatal maternal depression and anxiety on the processing of infant faces. *J Affect Disord* 2011;133(1–2):197–203.
- Dennis CL, McQueen K. The relationship between infant-feeding outcomes and postpartum depression: a qualitative systematic review. *Pediatrics* 2009;123:e73–e751.
- Halligan SL, Herbert J, Goodyer IM, *et al.* Exposure to postnatal depression predicts elevated cortisol in adolescent offspring. *Biol Psychiatry* 2004;55:376–81.
- Jones NA, Field T, Almeida A. Right frontal EEG asymmetry and behavioral inhibition in infants of depressed mothers. *Infant Behav Dev* 2009;32:298–304.
- Sohr-Preston SL, Scaramella LV. Implications of timing of maternal depressive symptoms for early cognitive and language development. *Clin Child Fam Psychol Rev* 2006;9:65–83.
- Thurgood S, Avery DM, Williamson L, *et al.* Postpartum depression (PPD). *American Journal of Clinical Medicine* 2009;6:17–22.
- Horowitz JA, Goodman JH. Identifying and treating postpartum depression. *J Obstet Gynecol Neonatal Nurs* 2005;34:264–73.
- Sokol LE, Epperson CN, Barber JP. Preventing postpartum depression: a meta-analytic review. *Clin Psychol Rev* 2013;33:1205–17.
- O'hara MW, Swain AM. Rates and risk of postpartum depression—a meta-analysis. *International Review of Psychiatry* 1996;8:37–54.
- Dennis CL, Chung-Lee L. Postpartum depression help-seeking barriers and maternal treatment preferences: a qualitative systematic review. *Birth* 2006;33:323–31.
- Gjerdingen DK, Yawn BP. Postpartum depression screening: importance, methods, barriers, and recommendations for practice. *J Am Board Fam Med* 2007;20:280–8.
- Cooney GM, Dwan K, Greig CA, *et al.* Exercise for depression. *Cochrane Database Syst Rev* 2013;9:CD004366.
- McCurdy AP, Boulé NG, Sivak A, *et al.* Effects of exercise on mild-to-moderate depressive symptoms in the postpartum period: a meta-analysis. *Obstet Gynecol* 2017;129:1087–97.
- Davies GA, Wolfe LA, Mottola MF, *et al.* Joint SOGC/CSEP clinical practice guideline: exercise in pregnancy and the postpartum period. *Can J Appl Physiol* 2003;28:329–41.
- American College of Sports Medicine. *ACSM's Guidelines for Exercise Testing and Prescription*. 9 edn. Philadelphia: Wolters Kluwer/Lippincott, Williams & Wilkins, 2013.
- ACOG Committee opinion. No 267, January 2002: exercise during pregnancy and the postpartum period. *Clin Obstet Gynecol* 2002;99:171–3.
- Mottola MF, Davenport MH, Ruchat S-M, *et al.* 2019 Canadian guideline for physical activity throughout pregnancy. *Br J Sports Med* 2018;52:1339–46.
- Moher D, Shamseer L, Clarke M, *et al.* Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015;4:1.
- ACOG Committee opinion no. 650: physical activity and exercise during pregnancy and the postpartum period. *Obstet Gynecol* 2015;126:e135–e142.
- Caspersen CJ, Powell KE, Christenson GM. Physical activity, exercise, and physical fitness: definitions and distinctions for health-related research. *Public Health Rep* 1985;100:126–31.
- Cochrane Collaboration. The Cochrane Collaboration's Tool for Assessing Risk of Bias, 2017. http://handbook.cochrane.org/chapter_8/8_5_the_cochrane_collaborations_tool_for_assessing_risk_of_bias.htm.

- 29 Guyatt GH, Oxman AD, Vist G, *et al*. GRADE guidelines: 4. Rating the quality of evidence—study limitations (risk of bias). *J Clin Epidemiol* 2011;64:407–15.
- 30 The Cochrane Collaboration. *Cochrane Handbook for Systematic Reviews of Interventions*. 5.1.0 edn, 2011.
- 31 Greenland S, Longnecker MP. Methods for trend estimation from summarized dose-response data, with applications to meta-analysis. *Am J Epidemiol* 1992;135:1301–9.
- 32 Orsini N, Li R, Wolk A, *et al*. Meta-analysis for linear and nonlinear dose-response relations: examples, an evaluation of approximations, and software. *Am J Epidemiol* 2012;175:66–73.
- 33 Viechtbauer W. Conducting Meta-Analyses in R with the metafor Package. *J Stat Softw* 2010;36:1–48.
- 34 R Core Team. *A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing, editor, 2017.
- 35 Harrell FE. *Regression Modeling Strategies with Applications to Linear Models, Logistic and Ordinal Regression and Survival Analysis*. 2nd edn. Cham: Springer-Verlag, 2015.
- 36 Satyapriya M, Nagarathna R, Padmalatha V, *et al*. Effect of integrated yoga on anxiety, depression & well being in normal pregnancy. *Complement Ther Clin Pract* 2013;19:230–6.
- 37 Gustafsson MK, Stafne SN, Romundstad PR, *et al*. The effects of an exercise programme during pregnancy on health-related quality of life in pregnant women: a Norwegian randomised controlled trial. *BJOG* 2016;123:1152–60.
- 38 Guskowska M, Langwald M, Sempolska K. Does physical exercise help maintain mental health during pregnancy? A comparison of changes in mental health in participants of physical exercise classes and childbirth classes. *J Phys Act Health* 2015;12:30–6.
- 39 El-Rafie MM, Khafagy GM, Gamal MG. Effect of aerobic exercise during pregnancy on antenatal depression. *Int J Womens Health* 2016;8:53–7.
- 40 Claesson IM, Klein S, Sydsjö G, *et al*. Physical activity and psychological well-being in obese pregnant and postpartum women attending a weight-gain restriction programme. *Midwifery* 2014;30:11–16.
- 41 Kalayeh S, Nikbakht R, Seddigi S. The effect of exercise during pregnancy on mother's depression in zahedan. *International Journal of Pharmacy and Technology* 2016;8:12248–62.
- 42 Downs DS, DiNallo JM, Kirner TL. Determinants of pregnancy and postpartum depression: prospective influences of depressive symptoms, body image satisfaction, and exercise behavior. *Ann Behav Med* 2008;36:54–63.
- 43 Loprinzi PD, Cardinal BJ, Loprinzi KL, *et al*. Benefits and environmental determinants of physical activity in children and adolescents. *Obes Facts* 2012;5:597–610.
- 44 Petrovic D, Perovic M, Lazovic B, *et al*. Association between walking, dysphoric mood and anxiety in late pregnancy: A cross-sectional study. *Psychiatry Res* 2016;246:360–3.
- 45 de Wit L, Jelsma JG, van Poppel MN, *et al*. Physical activity, depressed mood and pregnancy worries in European obese pregnant women: results from the DALI study. *BMC Pregnancy Childbirth* 2015;15:158.
- 46 Bershadsky S, Trumpfheller L, Kimble HB, *et al*. The effect of prenatal Hatha yoga on affect, cortisol and depressive symptoms. *Complement Ther Clin Pract* 2014;20:106–13.
- 47 Da Costa D, Rippen N, Dritsa M, *et al*. Self-reported leisure-time physical activity during pregnancy and relationship to psychological well-being. *J Psychosom Obstet Gynaecol* 2003;24:111–9.
- 48 Goodwin A, Astbury J, McMeeken J. Body image and psychological well-being in pregnancy. A comparison of exercisers and non-exercisers. *Aust N Z J Obstet Gynaecol* 2000;40:442–7.
- 49 Gaston A, Prapavessis H, Tired PH. Tired, moody and pregnant? Exercise may be the answer. *Psychol Health* 2013;28:1353–69.
- 50 Bowen A, Stewart N, Baetz M, *et al*. Antenatal depression in socially high-risk women in Canada. *J Epidemiol Community Health* 2009;63:414–6.
- 51 Battle CL, Uebelacker LA, Magee SR, *et al*. Potential for prenatal yoga to serve as an intervention to treat depression during pregnancy. *Womens Health Issues* 2015;25:134–41.
- 52 Ersek JL, Brunner Huber LR. Physical activity prior to and during pregnancy and risk of postpartum depressive symptoms. *J Obstet Gynecol Neonatal Nurs* 2009;38:556–66.
- 53 Ström M, Mortensen EL, Halldorson TI, *et al*. Leisure-time physical activity in pregnancy and risk of postpartum depression: a prospective study in a large national birth cohort. *J Clin Psychiatry* 2009;70:1707–14.
- 54 Beddoe AE, Paul Yang CP, Kennedy HP, *et al*. The effects of mindfulness-based yoga during pregnancy on maternal psychological and physical distress. *J Obstet Gynecol Neonatal Nurs* 2009;38:310–9.
- 55 Miquelutti MA, Cecatti JG, Makuch MY. Evaluation of a birth preparation program on lumbopelvic pain, urinary incontinence, anxiety and exercise: a randomized controlled trial. *BMC Pregnancy Childbirth* 2013;13:154.
- 56 Dodd JM, Newman A, Moran LJ, *et al*. The effect of antenatal dietary and lifestyle advice for women who are overweight or obese on emotional well-being: the LIMIT randomized trial. *Acta Obstet Gynecol Scand* 2016;95:309–18.
- 57 Newham JJ, Wittkowski A, Hurley J, *et al*. Effects of antenatal yoga on maternal anxiety and depression: a randomized controlled trial. *Depress Anxiety* 2014;31:631–40.
- 58 Shim MJ, Lee YS, Oh HE, He O, *et al*. Effects of a back-pain-reducing program during pregnancy for Korean women: a non-equivalent control-group pretest-posttest study. *Int J Nurs Stud* 2007;44:19–28.
- 59 Guskowska M. Physical activity in relation to affective states and labor anxiety in pregnant women. *Medicina Sportiva* 2011;15:114–8.
- 60 Field T, Diego M, Delgado J, *et al*. Yoga and social support reduce prenatal depression, anxiety and cortisol. *J Bodyw Mov Ther* 2013;17:397–403.
- 61 Ussher M, Lewis S, Aveyard P, *et al*. Physical activity for smoking cessation in pregnancy: randomised controlled trial. *BMJ* 2015;350:h2145.
- 62 Seime RJ, Vickers KS. The challenges of treating depression with exercise: from evidence to practice. *Clinical Psychology: Science and Practice* 2006;13:194–7.
- 63 Choi J, Fukuoka Y, Lee JH. The effects of physical activity and physical activity plus diet interventions on body weight in overweight or obese women who are pregnant or in postpartum: a systematic review and meta-analysis of randomized controlled trials. *Prev Med* 2013;56:351–64.
- 64 Rauff EL, Downs DS. Mediating effects of body image satisfaction on exercise behavior, depressive symptoms, and gestational weight gain in pregnancy. *Ann Behav Med* 2011;42:381–90.
- 65 Heyman E, Gamelin FX, Goekint M, *et al*. Intense exercise increases circulating endocannabinoid and BDNF levels in humans—possible implications for reward and depression. *Psychoneuroendocrinology* 2012;37:844–51.
- 66 Brändle E, Fritzsche G, Greven J. Affinity of different local anesthetic drugs and catecholamines for the contraluminal transport system for organic cations in proximal tubules of rat kidneys. *J Pharmacol Exp Ther* 1992;260:734–41.
- 67 Cramer H, Lauche R, Langhorst J, *et al*. Yoga for depression: a systematic review and meta-analysis. *Depress Anxiety* 2013;30:1068–83.
- 68 Syvälahti EK. Biological aspects of depression. *Acta Psychiatr Scand Suppl* 1994;377:11–15.

© 2018 Author(s) (or their employer(s)) 2018. No commercial re-use. See rights and permissions. Published by BMJ.